



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

decrease in size of the canines, and a development of an arched form of the lower jaws. He complains that many paleontologists have not appreciated the law of increase in size as a fundamental one in evolution, but, if the mammalogists have not appreciated it, surely other paleontologists have.

As regards the relationships between the South American and the Old World and North American mammalian faunas he says: "While the other orders are already represented in the South American *Notostylops* fauna, we have to deal, especially in the rodents and primates, with new faunal elements which must have gone thither either in the Oligocene or at the beginning of the Miocene. And they could have gone only from Europe or northern Africa, since, as we have seen, these rodents are closely related to the European forms, and the primates have at least a closer relationship with those of the Fayum than with those of the North American Eocene. There must, therefore, have been a connection between South America and the Old World in the Oligocene or at the beginning of the Miocene." This theory has already been urged by Ameghino. "This connection could not have been a broad land bridge, otherwise there would have been an exchange of the larger mammals, which did not occur till the Pliocene." He suggests that this migration of the smaller animals may have occurred from island to island of an archipelago, the creatures possibly carried by the larger birds of prey. And he thinks also that about the same time there was a like exchange of the smaller mammals between North America and Africa.

S. W. W.

---

*The Cid Mining District of Davidson County, North Carolina.* By JOSEPH E. POGUE. Raleigh: *Bull. No. 22 North Carolina Geol. Survey*, 1910. Pp. 144; Plates 22.

This district is located in the central portion of the Piedmont Plateau and includes areas of slate, tuffs, volcanic breccia, rhyolite, dacite, and andesite, cut by gabbro and diabase dikes. All but the dike rocks range from a massive to a schistose condition with sericite and greenstone schists as the final product of dynamic metamorphism. The slates are interbedded with rhyolitic and dacitic tuffs. The coarser acid volcanic breccia grades into rhyolite flows and is thought to be a flow breccia. The gabbro dikes are approximately parallel to the schistosity and are cut by diabase dikes, said to be Triassic. The evidence as to the Triassic

age of the latter is inconclusive as it rests on the fact that they cut Triassic sandstones. On this evidence they might be post-Triassic.

The region comprises a series of folds, beveled to the present surface, and one great overthrust fault. The jointing, folding, faulting, and schistosity are referred to the same epoch of compression.

Four types of ore bodies are noted, namely impregnations in the schists, stringer leads in quartz, parallel to the schistosity, quartz veins cutting the schistosity, and replacement deposits.

The ore minerals are auriferous pyrite, chalcopyrite, galena, and zinc blend. The deposits are referred to magmatic waters, perhaps emanating from a granitic mass a few miles west. The time of deposition is placed after metamorphism. In view of Emmons' recent work in Maine and Tennessee the evidence on this last point needs to be more carefully worked out.

A. D. B.

---

*The Iron Ore Supply of Japan.* By KINOSUKE INOUE. "The Iron Ore Resources of the World." Stockholm, 1910. Pp. 927-69; Plates 4; Figs. 13.

The iron ore deposits of Japan are classified in six groups as follows:

I. Magmatic segregations in granite. Not of economic importance under present conditions.

II. Bedded deposits usually in connection with radiolarian quartzites and slates of Paleozoic and Mesozoic age. The ores carry from 20 to 50 per cent iron with silica up to 40 per cent. They are usually rather high in phosphorus.

III. Contact deposits in limestone near contact with intrusives. These are the most important ores of Japan. The ore is chiefly magnetite with minor amounts of micaceous hematite and limonite. The iron content averages from 55 to 60 per cent with some analyses giving over 69 per cent. The ores are mixed with contact minerals and quartz and in some cases contain pyrite and chalcopyrite.

IV. Veins in various kinds of rocks. Not of great importance under present conditions.

V. Limonite deposits derived from the decomposition and redeposition of pyrite or magnetite deposits or by deposition from ferruginous springs. These are next in importance to class III.

VI. Alluvial deposits of iron sand derived from the decomposition of older rocks.

The amount of ore in sight is estimated at 19,000,000 metric tons;